



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A2	(11) International Publication Number: WO 98/03677 (43) International Publication Date: 29 January 1998 (29.01.98)
(21) International Application Number: PCT/EP97/03939 (22) International Filing Date: 22 July 1997 (22.07.97) (30) Priority Data: 96401634.9 22 July 1996 (22.07.96) EP <i>(34) Countries for which the regional or international application was filed:</i> GB et al. (71) Applicant (for all designated States except US): BYOCORE LIMITED [GB/-]; P.O. Box 3175, Road Town, Tortola (VG). (72) Inventors; and (75) Inventors/Applicants (for US only): RENAVILLE, Robert [BE/BE]; Rue de Moha 18, B-5030 Gembloux (BE). PORTETELLE, Daniel [BE/BE]; Rue de la Motte 11, B-5081 Meux la Bruyère (BE). (74) Agents: GUTMANN, Ernest et al.; Ernest Gutmann-Yves Plasseraud S.A., 3, rue Chauveau-Lagarde, F-75008 Paris (FR).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PIT-1 GENE POLYMORPHISM AND TRAIT SELECTION IN ANIMALS		
(57) Abstract <p>The invention concerns a genetic marker used to distinguish amongst animals a trait for milk producing capabilities or meat producing capabilities said genetic marker comprising a mutation in a fragment of a Pit-1 gene wherein three patterns of alleles are observed in which one of said allele patterns was fully mutated and being indicative of a trait of muscularity in said animal, while the two other allelic patterns, one being mutated and non/mutated, the other being non-mutated/non-mutated being indicative of a milk producing trait in said animal.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PIT-1 GENE POLYMORPHISM AND TRAIT SELECTION IN ANIMALSBACKGROUND OF THE INVENTION

5

Field of the Invention

The present invention relates to a genetic marker associated with different conformational traits. More specifically, the present invention describes a process wherein a polymorphism in a Pit-1 gene is used to determine traits in animals such as milk production and muscularity with ease.

10

Description of the Prior Art

Selection of a particular trait in a mammal is presently very expensive and very slow. Usually the selection process involves a geneological evaluation of the mammals history over a long period of time. This evaluation is based on various traits of the mammal or animal such as birth weight, growth weight, build, muscle strength, firmness, marbling, color, and the like.

15

Most of the selection of a particular trait in an animal to date, involves visually characterizing the specific traits over a time frame or weighing the animal at particular times. The animals with the quality traits that are to be selected are then bred with similar animals such that the particular trait is hopefully dominant in the next generation or the generations to follow.

20

The present methods for trait selection in mammals are often tedious and open to judgment of an expert in the field, such as a breeder. However, there is never any real assurance that the choice being made will dominate over the forthcoming generations. For example, in order to select a cow that is a good milk producing animal, it takes between 36 to 48 months to make such choice and after the choice is made, it is often based on hypothesis and the breeder's judgment.

25

30

In view of the uncertainty, expense and time involved with the current methods of trait selection in animals, new methods are currently under development which methods utilize a more scientific process which will hopefully improve the selection process.

5 One such method is the study of candidate genes to determine whether specific genes are associated with conformational traits in mammals and therefore these genes can be used as molecular markers to select particular traits of interest. This method first requires identification of candidate genes or anonymous genetic markers associated with the traits of
10 interest. The candidate gene approach can be successful, but first genes must be identified in the species of interest and correlated to the traits of interest.

The somatotropin system has several genes that may play a role in the control of particular traits in animals since this system is associated with
15 growth, lactation, reproduction and immunity. The somatotropin system is quite complicated and involves at a hypothalamic level, somatocinin and somatostatin; at a pituitary level, pituitary-specific transcription factor (Pit-1) which is responsible for growth hormone expression in mammals; at a hepatic level, growth hormone receptor and growth hormone plasmatic transport
20 protein; and at a cellular level, growth hormone receptor, insulin-growth factor-1 and insulin growth factor transport protein.

Selection of genes from this somatotropin system that may influence particular traits in animals is quite complicated, since this system has many different functions in different parts of the animal, from the pituitary to the
25 cellular level.

The present invention involves the selection of a gene, the pituitary-specific transcription factor (hereinafter referred to as Pit-1) that can act as a genetic marker to characterize specific traits in animals.

Pit-1 is a member of the POU family of homeo-domain transcription
30 factors and plays an important role in developmental processes. The POU-domain was originally identified as a highly conserved region of 150 to 160

amino acids found in three mammalian transcription factors, Pit-1, Oct-1, Oct-2 and also in the product of nematode gene *unc-86* (Herr et al., *Genes & Dev.* **2**: 1513 (1988); Ruvkun and Finney, *Cell* **64**:475 (1991)).

Pit-1 is a pituitary-specific transcription factor that regulates growth hormone, activates prolactin and has a role in pituitary cell differentiation and proliferation (Steinfelder et al., *P.N.A.S., USA* **88**:3130 (1991). Mutations in the Pit-1 gene responsible for the dwarf phenotypes of the Snell and Jackson mice and lead to anterior pituitary hypoplasia (Li et al., *Nature* **347**:528 (1992)). Moreover, it has been shown that the inhibition of Pit-1 synthesis leads to a decrease in prolactin and growth hormone (GH) expression and to a dramatic decrease in cell proliferation in GH and prolactin producing cell lines (McCormick et al., *Nature* **345**:829 (1990)).

In human, different mutations in the Pit-1 gene have also been reported in patients with familial pituitary hypoplasia (Pfaffle et al., *Science* **257**:1118 (1992)); and in patients with sporadic combined pituitary hormone deficiency (Radovick et al., *Science* **257**:1115 (1992); Tatsumi et al., *Nature Genetics* **1**: 56 (1992)).

The cDNA sequence of bovine Pit-1 has been published by Bodner, M. et al, *Cell* **55** (3): 505-568 (1988) and is shown in Figure 2.

The Association of Pit-1 polymorphisms with growth and carcass traits in pigs has been described by Yu et al., *J. Anim. Sci.* **73**: 1282 (1995). Yu et al., *supra* described three Pit-1 polymorphisms in pigs based on two restriction fragment length polymorphisms (hereinafter referred to as RFLP) using a Pit-1 POU-domain cDNA probe and the restriction enzymes *Bam*HI and *Msp*I and a PCR/RFLP using *Rsa*I.

Results from Yu et al.'s, *supra*, mixed-model analysis revealed that pigs with the *Msp*I CC genotype were associated with heavier birth rate than the DD genotype pigs. Moreover, with the Pit-1 *Bam*HI polymorphisms heavier birth weight was significantly associated with the BB genotype, although the authors cautioned against concluding such association since the BB genotype population was extremely small.

Although Woolard et al., *J. Anim. Sci.* 72:3267 1994) recognized a HinfI polymorphism at the bovine Pit-1 gene locus, these authors failed to link this mutation to the selection trait in animals. The conclusion drawn in Woolard, *supra* was that polymorphic fragments that were observed were
5 consistent with autosomal Mendelian inheritance.

There is no disclosure in Yu et al. or Woolard et al of any association of the allele pattern AB with milk production, nor the allele pattern BB with muscularity in animals.

Therefore, the present invention overcomes the disadvantages of the
10 current methods of trait selection in animals by providing a scientific basis for selection of traits by use of a genetic marker.

Moreover, the process described in the present invention can be used to characterize superior milk producing animals from animals having meat producing characteristics.

15 It has been surprisingly discovered that a polymorphism in the Pit-1 gene can be used to characterize traits such as milk production and muscularity in animals. Two alleles, A and B were distinguished for the Pit-1 gene responsible for the activation of prolactin and growth hormone gene expression. The AA pattern was less frequent than the AB or BB pattern. The
20 significant superiority of the Pit-1 AB pattern or AA pattern over the BB pattern was observed for milk, protein and angularity. Likewise the BB genotype pattern was associated with animal muscularity.

This discovery permits the use of the mutation in the Pit-1 gene to be utilized as a genetic marker to identify certain traits in animals. Once these
25 particular traits are identified, the animals can be sold at market with increased value due to their superior traits.

Accordingly, it is an object of the present invention to provide a genetic marker for trait selection in animals.

In another aspect, the present invention provides a process to
30 characterize animals having superior milk production traits or muscularity traits.

In yet another aspect, the present invention provides genetically engineered animals that have superior milk production, angularity, fat, protein or muscularity traits. These and other objects are achieved by the present invention as evidenced by the summary of the invention, description
5 of the preferred embodiments and the claims.

SUMMARY OF THE INVENTION

The present invention thus provides a genetic marker that can be used for trait selection in mammals.

Furthermore, the present invention provides a method to identify a
10 polymorphism present in the Pit-1 gene which polymorphism can be utilized to select superior traits in animals for angularity, fat, muscularity, protein or milk production.

Accordingly, in one of the composition aspects, the present invention relates to a genetic marker used to distinguish amongst animals a trait for
15 milk producing capabilities or meat producing capabilities said genetic marker comprising a mutation in a fragment of a Pit-1 gene, wherein three allele patterns are observed, the fully mutated pattern being indicative.

In the present application, the marker characteristic of milk producing capabilities is called AA for its homozygous state of the allele and the marker
20 characteristic of meat producing capabilities is called BB for its homozygous state.

The sequences of alleles A and B differ only by one transition from the Adenosine in position 1178 of the sequence of Figure 2 in Pit-1 AA to a
25 guanine in Pit-1 BB, as demonstrated by the inventors by experiments shown in Example B.

In a preferred embodiment for the present invention, the three allele patterns are distinguished after digestion with a restriction endonuclease, which cleaves the mutated Pit-1 gene fragment and not the non-mutated Pit-1
30 gene fragment, the fully digested pattern being indicative of a trait for muscularity in said animal, while the intermediate digested/non-digested

pattern or the fully non-digested pattern being indicative of a milk producing trait in said animal.

In a more preferred embodiment of the present invention the restriction endonuclease utilized is *Hinfl*.

5 In another preferred embodiment of the present invention, the three allele patterns are distinguished using probes which overlap the mutated region in said Pit-1 gene, one probe being specific for the mutated Pit-1 gene and another one being specific for the non-mutated Pit-1 gene.

In another aspect, the present invention relates to a process for
10 detecting certain traits in an animal, said process comprising the steps of:

(1) isolating genomic DNA from an animal;

(2) optionally isolating a fragment from said genomic DNA comprising a fragment of a Pit-1 gene;

(3) detecting a mutation in the Pit-1 gene; and

15 (4) analyzing said mutation to determine a trait in said animal wherein upon analysis traits of muscularity and fat can be distinguished from milk producing traits in said animals.

In particular embodiment of the present invention, detection is accomplished by using restriction endonucleases.

20 In another particular embodiment of the present invention, detection is accomplished by using probes which overlap the mutated gene in said Pit-1 gene, more particularly the 1178 position.

In yet another aspect, the present invention relates to genetically engineered animals that have the characteristic traits described in the present
25 invention.

BRIEF DESCRIPTION OF THE DRAWINGS

30 Fig. 1 is an electrophoretic gel illustrating the PCR/Restriction Fragment Length Polymorphism patterns using the restriction enzymes *Hinfl* on the Pit-1 gene observed in Holstein-Friesian and Simmental Bulls. The

sizes of digested fragments are on the left, and the patterns are at the top. Fragment length (in kilobases) was estimated relative to the DNA size markers ϕ X174 DNA/HaeIII fragments.

Figure 2 is the sequence of bovine Pit-1 cDNA.

5 Figure 3 is an electrophoretic pattern illustrating the PCR amplification products obtained after amplification with following primers:

lines 1-3-5 : Pit 1 AA and Pit 1 B,

lines 2-4-6 : Pit 1 BB and Pit 1 B.

10 DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS
OF THE INVENTION

As used herein, the "animal" encompasses all mammals, avians, and fish including but not limited to, cows, bulls, goats, pigs, sheep, chickens and
15 the like. In view of the high degree of conservation of the Pit-1 gene among species (> 95 %) the invention is easily transposable from one specie to another. Also, the instant invention can be used in human beings to determine traits such as capacity to metabolize growth hormone.

The term "polymorphism" refers to the simultaneous occurrence in the
20 population of genomes showing allelic variations as seen either in alleles producing different phenotypes or in changes in DNA affecting the restriction pattern.

As used herein the term "trait" encompasses any characteristic, especially one that distinguishes one animal from another.

25 The term "angularity", as used herein means an objective criteria used to identify specific traits of an animal in relation to specific measurements which can be taken on the animal's body. The measurements are taken on the animal with respect to certain morphological characteristics.

For example, to determine the angularity for a milk production trait, the
30 pelvic bones and muscles surrounding the pelvic bones of an animal are measured to determine whether they are projecting or not. A scale can then

be established. When the bones are very projecting, there are very little rounded muscles and thus the animals are milk producing. To the contrary, when the bones are not protruding and there are a lot of rounded muscle present on the animal, the animal would not be considered a good milk produce, but rather a beef producer.

As used herein, the term "muscularity" encompasses animals that are better meat producers that can be slaughtered for their meat than milk producers.

More specifically, the present invention relates to the use of a Pit-1 gene polymorphism as a potential marker for genetic variations in animals. Pit-1 codes for a factor of transcription in a cell and any mutation of this gene can alter by diminution or augmentation the capacity of transcription thus resulting in polymorphisms which effect the outcome of different traits in an animal.

The Pit-1 gene was previously identified in a 13-kb bovine genomic library by Woolard et al., *supra*. A 13-kb clone was isolated from this library by using a bovine Pit-1 cDNA, which is labeled, as a probe of:

5'-AAACCATCATCTCCCTTCTT-3' (SEQ. ID. NO: 1)

5'- AATGTACAATGTGCCTTCTGAG-3' (SEQ. ID. NO:2).

Characterization of *Xho*I, *Hin*fI and *Eco*RI subclones of this 13-kb insert by restriction enzyme digestion and sequencing identified this clone as a bovine Pit-1 genomic fragment.

Similarly methods as taught by Woolard et al, *supra* can be used to identify the Pit-1 gene in different genomic libraries other than bovine. This will permit the identification of specific sequences within the Pit-1 genomic fragment that can be used to amplify this sequence from different animals as described below.

The first step in identifying a mutation in the Pit-1 gene in an animal is to obtain a sample from the animal such as, but not limited to semen, blood,

cells, biopsy tissues, feces and the like. Genomic DNA can then be extracted for the specimens obtained using methods known in the art as described by Sambrook et al., *Molecular Cloning, A Laboratory Manual*, second edition 1989.

5 However, it is preferable to extract the genomic DNA using the procedure described in Walsh, *Biotechniques*, **10**:506 (1991) for semen or the procedure for blood as described by Lewin and Stewart-Haynes *Biotechniques*, **13**:522.

10 After extracting the genomic DNA there are several known methods in the art to detect the mutation in the Pit-1 gene. Any detection method can be utilized to detect the mutation. Examples of these methods include, but are not limited to RFLP, SSCP, DGGE, CFLP and single base mutations as described by Prosser, *Trends Biotech* **11**:238-246 (1993) and Sambrook et al., *supra*. These methods will be discussed in greater detail below.

15 For example, in the RFLP (restriction fragment length polymorphism) method, PCR primers are used to amplify by standard procedures a fragment that includes the Pit-1 gene. Any PCR primers can be utilized that would permit the amplification of the Pit-1 sequence and the method in isolating the particular clone which would identify such primers.

20 In a preferred embodiment of the invention, the PCR primers can be designed from intron V and exon 6 of a fragment containing the polymorphism of the Pit-1 gene, such as the 451-bp fragment described by Woolard et al., *supra*. In a more preferred embodiment of the present invention, the PCR primers are as follows:

25

5'-AAACCATCATCTCCCTTCTT-3' (SEQ. ID. NO: 1)

5'- AATGTACAATGTGCCTTCTGAG-3' (SEQ. ID. NO:2)

30 Amplification of the Pit-1 fragment can be performed using standard PCR procedures, as described in Sambrook et al., *supra*. It is preferable,

however, to amplify the genomic DNA in a 50 μ l reaction volumes containing 2 mM MgCl₂.

In a preferred embodiment of the invention, the following conditions for the PCR reaction can be employed: between 88°C to 98°C for 10 to 15 minutes; and between 90°C to 100°C for about 1 minute, followed by between 25 to 50 cycles at between 90°C to 100°C for 20 to 40 seconds; 40°C to 60°C for 1 to 5 minutes; and 68°C to 80°C for about 1 to 5 minutes. The last step may encompass a cycle at between 68°C to 80°C for 8 to 12 minutes.

After amplification the particular mutation in Pit-1 is then cut using various restriction enzymes or endonucleases known in the art. These restriction enzymes include, but are not limited to *Bam*HI, *Eco*RI, *Sma*I, *Hin*fl and the like. See, for example those enzymes described in Sambrook et al., *supra*. It is of particular interest to use a restriction endonuclease which cleaves the mutated allele of the Pit-1 gene and does not cleave the non-mutated allele of the Pit-1 gene. In a preferred embodiment of the present invention with the respect to the identification of milk production, fat, protein and muscularity traits in animals, *Hin*fl is utilized.

After digestion, the sample is then electrophoresed on agarose gels and identified with a stain such as, for example ethidium bromide, however any stain can be used that identifies the fragments

SCCP(single stranded conformation polymorphism) is also a method known in the art that can identify a mutation or mutations in the isolated genomic Pit-1 fragment. This method is based on PCR amplification, using similar primers as those described above. The amplified fragment is then labeled with a label such as ³²P or with any other appropriate radioactive label. The radiolabeled fragment is then denatured, for example by heating and then subjected to quick cooling. After cooling, the fragment is then electrophoresed using non-denatured technique and then autoradiographed.

DGGE (denaturing gradient gel electrophoresis) is yet another method to detect the Pit-1 mutation. In this process, the fragment is amplified by PCR using appropriate primers, such as those described above and subjected to a

denaturing gradient. The sample is further electrophoresed and the mutation is detected.

Yet another method that can be used to detect the mutation is CFLP (cleavage fragment length polymorphism). This method can detect mutations of a sole base in the DNA sequence between two molecules of wild-type DNA and of a mutant type of DNA. This method is now marketed by Boehringer Mannheim and can be purchased in the form of a kit.

Another method that can be used to detect the mutation in the Pit-1 gene utilizes primers that overlap the mutated region of the Pit-1 gene.

More preferably, two separate amplification reactions are performed on the extracted genomic DNA sample using two sets of primers, one set containing a primer which overlaps and is specific for the mutated Pit-1 gene, another set containing a primer which overlaps and is specific for the non-mutated Pit-1 gene. Even more preferably, the primers used are labeled so that the amplification product can be easily visualized. According to this method, when the tested genomic DNA contains a homozygous mutated Pit-1 gene (two mutated alleles), only the amplification reaction using the probe specific for the mutated region will produce a signal (i.e., an amplification product). When the tested genomic DNA is heterozygous (one mutated allele and one non-mutated allele) the two amplification reactions will produce a signal (an amplification product). Similarly, when the tested genomic DNA contains a homozygous non-mutated Pit-1 gene, only the amplification reaction using the probe specific for the non-mutated region will produce a signal. Therefore, in one single amplification step, the allele pattern of the tested DNA becomes apparent.

In a second embodiment, the use of technique described in WO 97/06276 is particularly adapted to detect in a single step the homozygous or heterozygous state of the marker.

In these methods, there is no need to further cut the amplified product in order to distinguish among the various patterns. Furthermore, there is no need to amplify a specific Pit-1 gene fragment prior to the detection step.

Finally, depending on the nature of the label used, the visualization can be very easy. For instance, when the probes are radiolabelled visualization is obtained by electrophoresis. More interestingly, when the probes are labelled with stains, immediate visualization is obtained.

5 In the first embodiment, the test can be carried out in very simple devices, such as plates. Samples of the genomic DNA are introduced into 2 wells, one with the labeled set of probes containing one probe which overlaps the Pit-1 gene mutation and is specific for the mutated Pit-1 gene, one with the set of probes containing one probe which overlaps the Pit-1 gene
10 mutation and is specific for the non-mutated Pit-1 gene.

After amplification, the labelling appears directly in the plates and can be analyzed by automated devices.

In this amplification method, the second primer used in each of the set of probes is selected in such a way as to enable amplification of a product
15 containing from 200 to 400 bp. More preferably 320-370 bp. Such second primers can be for instance selected from the following primers:

gac agggaaagtg atatagaaag ggagataga (Pit-1 B)

The length of each of the primers is preferably comprised between 20
20 and 40 bases, more preferably between 25 and 35.

The selection of the appropriate probes for this strategy has been made possible by the identification, by the inventors, of a mutation in the Pit-1 gene that is responsible for the observed polymorphism. More specifically, this mutation occurs in the Pit-1 coding region, at nucleotide 1178, where an
25 Adenine is substituted in the mutated gene by a guanine. This mutation is shown in Figure 2.

The position of the probe which overlap the mutated region (the mutation) can vary.

More preferably, in the first embodiment, the two couples of primers
30 are:

ca gagagaaaaa cgggtgaaga caagcat a (Pit-1 AA)

gac agggaaagtg atatagaaag ggagataga (Pit-1 B)
for the AA genotype, characteristic of milk producing capabilities. and

ca gagagaaaaa cgggtgaaga caagcat g (Pit-1 BB)

5 gac agggaaagtg atatagaaag ggagataga (Pit-1 B)
for the BB genotype, characteristic of meat producing capabilities.

In the second embodiment, when the method of WO 97/06276 is used,
two couples of primers leading to the production of amplified fragments of
10 different sizes are:

ca gagagaaaaa cgggtgaaga caagcat a (Pit-1 AA)

gac agggaaagtg atatagaaag ggagataga (Pit-1 B)

for the AA genotype, characteristic of milk producing capabilities, and

15 ca gagagaaaaa cgggtgaaga caagcat g (Pit-1 BB)

and a second primer Pit-1 C' chosen in such a way that the amplification is at
least 10 bp shorter or longer than those obtained with Pit-1 AA and Pit-1B for
the BB genotype, characteristic of meat producing capabilities.

Although many detection methods for mutations are available, the
20 present invention is not limited to the methods discussed above and
encompasses all methods for detecting a mutation.

The alleles and allelic patterns are then identified and statistical
analysis is then performed to determine the specific traits evidenced by the
identification of the alleles. More specifically, any statistical program that can
25 identify daughter yield variations (DYD) and deregressed proofs (DRP) can
be utilized. It is preferable to perform the statistical analysis using the
MIXED procedure of SAS (User's Guide:Statistics, Version 6, 4th ed. SAS
Inst., Inc. Cary, N.C. (1990); Technical Report P 229 SAS Inst., Inc., Cary,
N.C. (1992). The statistical analysis used in the present invention is
30 discussed in detail in the examples below.

Also encompassed by the present invention is a kit containing extraction materials for genomic DNA, the PCR primers having SEQ ID NOS. 1 and 2 (illustrated above), the materials necessary to visualize the mutation such as electrophoretic gels and the like. The content of the kit may vary depending upon the detection methods utilized, which are discussed in detail above.

Also encompassed by the present invention are primers that overlap the mutation in the Pit-1 gene.

More specifically, the invention also relates to a primer comprising from 20 to 40 bases, which is complementary to a region of the Pit-1 gene having a mutation.

The invention also embraces sets of primers which allow the amplification of a region of 200 to 400 bases in the Pit-1 gene, wherein said region contains a mutation.

The following primers are encompassed in the present invention:

ca gagagaaaaa cgggtgaaga caagcat a (Pit-1 AA)

ca gagagaaaaa cgggtgaaga caagcat g (Pit-1 BB)

gac agggaaagtg atatagaaag ggagataga (Pit-1 B)

In order to further illustrate the present invention and advantages thereof, the following specific examples are given, it being understood that the same are intended only as illustrative and in nowise limitative.

EXAMPLE A

25

1. DNA EXTRACTION AND PCR

Genomic DNA of 89 commercially available registered Italian Holstein-Friesian bulls was extracted from semen as described by Lucy et al., *Domest. Anim. Endocrinol.* 10:325 (1993).

The RFLP at the Pit-1 gene using *HinfI* restriction enzyme was revealed by PCR analysis adapted from Woolard et al., *supra*.

The PCR primers were designed from intron V and exon 6. The sequences of the primers used were 5'-AAACCATCATCTCCCTTCTT-3' (SEQ ID NO:1) and 5'-AATGTACAATGTGCCTTCTGAG-3' (SEQ ID NO:2). These primers were used to amplify by standard procedures a 451-bp fragment from the genomic DNA in 50- μ L reaction volumes containing 2 mM MgCl₂. Conditions were 94.5°C, 10 min., and 94°C, 1 min., followed by 35 cycles of 95°C, 30 s, 56°C, 1 min., and 72°C, 2 min. The last step was 72°C for 10 min. PCR products were digested with *Hinf*I and electrophoresed on 2% agarose gels with 1 μ g/mL ethidium bromide (Figure 1).

Daughter yield deviations (DYD) computed in March 1996, were obtained from the Holstein-Friesian bulls from the Italian Holstein-Friesian Breeder Association ANAFI (Associazione Nazionale Allevatori Frison Italiana, Cremona, Italy). DYD values are not computed for fat and protein percentage as those traits are only evaluated indirectly out of solutions for yield traits and mean population values for those traits. Therefore DYD values were computed using the same approach as for the computation of genetic values for percentage traits.

Similar DYD were also not available for type traits, therefore genetic values were transformed to deregressed proofs (DRP) (Banos et al., Interbull Annual Meeting, Aarhus, Denmark, Bulletin No. 8, 1993, Sigbjorn et al., *J. Dairy Sci.*, 78:2047 (1995) that can then be considered approximate DYD.

Means and standard deviations of DYD for milk production traits and or DRP for conformation traits of the bulls sample are presented in Table I. Effective number of daughters, which is a measure of the number of daughters adjusted for their distribution inside herds was available for yield traits, but not for type traits. It was therefore approximated using the following formula: effective number = real number x square root of ratio between number of herds and number of daughters.

TABLE I

TABLE 1. Mean daughter yield deviations for milk traits and deregressed proofs for conformation traits of 89 Holstein-Friesian bulls.

Trait	\bar{X}	SD	Minimum	Maximum
Milk traits				
Milk, kg	+317	221	-231	+899
Fat, kg	+10.8	8.2	-14	+28
Protein, kg	+11.6	7.2	-7	+32
Fat, % ¹	-0.003	0.091	-0.17	+0.23
Protein, % ¹	+0.021	0.045	-0.11	+0.12
Effective daughters ²	490	1443	69	10298
Conformation traits³				
Final score	+0.147	0.438	-0.75	+1.19
Stature	+0.210	1.536	-3.76	+4.56
Strength	+0.218	1.662	-3.68	+3.46
Body depth	+0.340	1.599	-3.42	+3.64
Angularity	+0.681	1.215	-3.44	+3.42
Rump angle	-0.111	1.807	-4.44	+4.10
Rump width	+0.007	1.591	-3.28	+4.34
Rear legs	+0.203	2.266	-5.66	+5.66
Fect	+0.053	1.746	-5.26	+3.84
Fore udder	+0.038	2.207	-5.44	+5.46
Height rear udder	+0.458	1.856	-3.64	+4.44
Width rear udder	+0.864	1.474	-2.80	+4.18
Udder support	+0.514	2.453	-10.72	+7.12
Udder depth	-0.282	1.702	-5.78	+3.74
Teat placement	+0.479	1.633	-4.12	+3.82
Teat length	+0.416	2.112	-4.60	+6.74
Effective daughters ⁴	195	471	18	3199

¹ Percentage fat and protein daughter yield deviations computed from yields.

² Number of effective daughters for yield reported by ANAFI.

³ Deregressed proofs for final score reported on original scale, for linear scores on relative scale.

⁴ Approximate number of effective daughters obtained from numbers of daughters and herds.

2. STATISTICAL ANALYSIS

Statistical analysis was performed using the MIXED procedure of SAS *supra*. The mixed model used was

$$y = Xb + Zu + e$$

Where

y = vector of DYD or DRP of bulls; b = vector of fixed effects associated with Pit-1 pattern, u = vector of random additive polygenic effect of bulls, and e = vector of random residual effects. This model was solved using the following mixed model equations:

$$\begin{bmatrix} X'R^{-1}X & X'R^{-1}Z \\ Z'R^{-1}X & Z'R^{-1}Z + A^{-1}\left(\frac{4-h^2}{h^2\hat{\sigma}_e^2}\right) \end{bmatrix} \begin{bmatrix} \hat{b} \\ \hat{u} \end{bmatrix} = \begin{bmatrix} X'R^{-1}y \\ Z'R^{-1}y \end{bmatrix} \Leftrightarrow \begin{bmatrix} C_{bb} & C_{bu} \\ C_{ub} & C_{uu} \end{bmatrix} \begin{bmatrix} \hat{b} \\ \hat{u} \end{bmatrix} = \begin{bmatrix} X'R^{-1}y \\ Z'R^{-1}y \end{bmatrix}$$

where A is the additive relationship matrix between the 89 bulls constructed using all known relationships (1842 known ancestors), $R^{-1} = D / \hat{\sigma}_e^2$, where D is assumed to be a diagonal matrix with the number of effective daughters for every bull on its diagonal. This matrix is then divided by the estimate of the residual variance $\hat{\sigma}_e^2$. This is a REML estimated (Patterson and Thompson, *Biometrika* 58:545 91971), here identical to non-interactive minimum variance quadratic unbiased estimation (Rao, *J. Mult. Anal.* 1:445 (1971), as convergence occurs after 1 round. The estimate found has the property of being the quadratic forms minimizing the sampling variance. Two assumption were made, no residual covariances between DYD or DRP and heritabilities (h^2) of DYD or DRP equal to heritabilities use for genetic evaluations with the exception of percentage of fat and protein where 0.50 was assumed to be the heritability (Table 2). This method tends to

overestimate additive heritability as variance due to sires is not reduce for the presence of the Pit-1 pattern in the model, but this overestimation should be not very important.

5

TABLE II

TABLE 2. Assumed heritabilities and milk traits and conformation traits of Italian Holsteins.

Trait	Heritability
Milk traits	
Milk, kg	0.25
Fat, kg	0.25
Protein, kg	0.25
Fat, % ¹	0.50
Protein, % ¹	0.50
Conformation traits	
Final score	0.15
Stature	0.38
Strength	0.29
Body depth	0.31
Angularity	0.31
Rump angle	0.25
Rump width	0.29
Rear legs	0.16
Feet	0.18
Fore udder	0.15
Height rear udder	0.20
Width rear udder	0.24
Udder support	0.15
Udder depth	0.29
Teat placement	0.22
Teat length	0.22

¹ Percentage fat and protein daughter yield deviations computed from yields, therefore assumed heritability is not the heritability used for breeding value estimation.

Linear contrasts were constructed as differences between pattern solutions. Testing of contrasts was done using the following statistic:

$$5 \quad F = b' l' l' C^{bb} l)^{-1} l' b$$

where $l'b$ represents differences between pattern solutions, l being the linear contrast vector, C^{bb} an estimate of the block of the generalized inverse of the coefficient matrix associated with pattern effects and $(l' C^{bb} l)^{-1}$ is the
 10 inverse of the squared standard error of the linear contrast. the numerator degree of freedom was approximated using $\text{rank}(l) = 1$. The denominator was put to $n - \text{rank}(X) = 86$ where n is the number of observations.

It is not certain that the presence of a given pattern has only one major effect. Therefore the following strategy based of Weller et al., *J. Dairy Sci.*
 15 **73**: 2525 (1990) was used to test this hypothesis.

1. Traits showing single-trait significant contrasts between patterns were grouped, eventual related traits were also included.
2. Weighted correlation V and covariance P matrixes among these traits were
 20 obtained.
3. A canonical transformation was defined as $V = QEQ'$, where E is a diagonal matrix of eigenvalues, and Q a matrix of eigenvectors.
4. The transformation matrix T was defined as $Q^{-1}S$ where S is a diagonal matrix of the inverse standard-deviations of the original traits, therefore
 25 $TPT' = E$.
5. The transformation matrix was used to transform the related traits to unrelated canonical traits.
6. Approximate heritabilities and weights for the canonical traits were obtained as weighted averages of the values for the initial traits. weighting
 30 coefficients were the squared values of Q^{-1} .

7. Canonical traits were analyzed using the methods described above for initial traits. Canonical traits showing only low relative eigenvalues explain little of the observed variance.

8. Multiple-trait linear contrasts for original effects can be estimated using
5 back transformation of significant canonical contrasts.

9. The results for these new traits are then useful to determine if only one effect of the Pit-1 pattern can be observed, or if there are more than one significant effects. Backtransformed contrasts reflect the significant differences between original traits based on a given effect of Pit-1 on the
10 canonical trait.

3. RESULTS

PCR/RFLP

15 The PCR product was 451 bp in length. Digestion of the PCR product with *HinfI* revealed two alleles: the A allele not digested with *HinfI* and yielding a 451 bp fragment and the B allele cut at one restriction site and generating two fragments of 244 and 207 bp in length as described by Woollard et al., *supra* (Figure 1).

20

Relationship of PCR/RFLP to Milk production

The frequencies of the three pattern AA, AB, and BB were 2.2%, 31.5% and 66.3%. The frequencies of the A and B alleles were estimated by a
25 maximum likelihood approach with 18.8% for A and 81.2% for B.

Table 3 shows the linear contrasts and standard errors between the three Pit-1 pattern. Therefore the highly significant contrasts ($P < 0.01$) observed for rear legs seem to be more due to the fact that the typed AA animals are extreme on this trait than to a real biological reason. Highly
30 significant contrasts between AB and BB patterns were found for milk and protein yield ($P < 0.01$). Significant contrasts were observed for fat

percentage and angularity ($P < 0.05$). The AB pattern or AA pattern was superior for milk, protein yield and angularity and inferior for fat percentage. These results can be interpreted as resulting from a single positive action of the heterozygote AB or AA on milk yield, thereby influencing protein yield positively and not fat yield which gives the observed negative influence on fat percentage. The influence of Pit-1 on angularity is in this context not very surprising as this linear trait is considered being strongly related to milk yield.

TABLE III

TABLE 3. Linear contrasts (C) and standard errors (SE) between the three Pit-1 patterns observed on 89 Holstein-Friesian bulls.

Trait	Contrast					
	AA-AB ¹		AA-BB ¹		AB-BB	
	C	SE	C	SE	C	SE
Milk traits						
Milk, kg	-152	156	-21	150	131 **	49
Fat, kg	5.0	5.7	5.4	5.7	0.4	1.8
Protein, kg	-4.2	4.9	0.8	4.5	4.9 **	1.5
Fat, % ²	0.114	0.062	0.067	0.062	-0.047 *	0.019
Protein, % ²	0.005	0.034	0.015	0.031	0.010	0.010
Conformation traits³						
Final score	-0.376	0.299	-0.253	0.298	0.123	0.092
Stature	-0.745	1.043	-0.501	1.044	0.244	0.329
Strength	0.915	1.143	1.012	1.138	0.097	0.367
Body depth	0.108	1.076	0.562	1.061	0.454	0.332
Angularity	-0.478	0.809	0.072	0.716	0.550	0.252 *
Rump angle	-0.211	1.219	-0.514	1.286	-0.303	0.398
Rump width	0.019	0.608	0.147	1.039	0.128	0.330
Rear legs	-4.404 ** ¹	1.548	-4.784 ** ¹	1.542	-0.380	0.479
Fect	1.588	1.264	1.731	1.259	0.142	0.395
Fore udder	-0.653	1.540	-1.256	1.546	-0.603	0.478
Height rear udder	-0.974	1.290	-0.998	1.288	-0.024	0.750
Width rear udder	-0.378	1.047	0.072	2.273	0.449	0.324
Udder support	-1.798	1.707	-1.157	1.706	0.641	0.525
Udder depth	-1.447	1.245	-1.673	1.240	-0.226	0.388
Teat placement	-1.385	1.158	-1.548	1.154	-0.163	0.356
Teat length	0.041	1.297	0.312	1.396	0.271	0.446

¹ Only 2.2 % of the animal were AA, therefore all results comparing this pattern are preliminary

² Percentage fat and protein daughter yield deviations computed from yields.

³ Deregressed proofs for final score reported on original scale, for linear scores on relative scale.

* P < 0.05

** P < 0.01

In order to test the hypothesis of a single action we performed a canonical transformation of milk, fat and protein yields. Yields were analyzed as percentage DYD were obtained as functions of yields; therefore this results in no new information. Angularity was added. The phenotypic correlation matrix was computed. Observations were weighted using the number of effective daughters. Since these numbers were different for yield and type traits approximate weights were obtained as weighted means of numbers of effective daughters. Table 4 gives the correlations. Correlations among yield traits showed the expected values with higher correlations between milk and protein than between fat and one of the other traits. Angularity showed correlations between 0.42 and 0.51 with yields traits.

TABLE 4. Correlations among daughter yield deviations for the milk traits and angularity.

Trait	Trait			
	Milk yield	Fat yield	Protein yield	Angularity
Milk yield	1.00	0.72	0.90	0.42
Fat yield		1.00	0.76	0.51
Protein yield			1.00	0.48
Angularity				1.00

Results from the canonical decomposition of the correlation matrix are in table V. The first and the second canonical trait explain 90% of the total variance. Especially the last canonical trait was not very informative. Table 5 gives also the eigenvectors and the relative importance of the different traits in each eigenvector. The first canonical trait is a combination of all four traits with relative influences between 15% for angularity and 30% for protein. The second canonical trait however is more specifically linked to angularity with a

relative importance of 81% in this trait. The third is associated with fat and less with milk, the fourth only with milk and protein.

TABLE 5. Standardized eigenvectors and eigenvalues of the four canonical traits (between brackets relative importance of eigenvalues in total variance and of values in eigenvectors in canonical traits).

Canonical Trait	Eigenvalue	Eigenvector			
		Milk yield	Fat yield	Protein yield	Angularity
1	2.94 (73%)	0.532 (28%)	0.515 (27%)	0.548 (30%)	0.389 (15%)
2	0.67 (17%)	0.349 (12%)	0.047 (<1%)	0.257 (7%)	-0.900 (81%)
3	0.30 (8%)	0.396 (16%)	-0.853 (73%)	0.283 (8%)	0.189 (4%)
4	0.09 (2%)	0.662 (44%)	0.072 (<1%)	-0.744 (55%)	0.048 (<1%)

Table 6 shows the linear contrasts and standard-errors observed for the four canonical traits. Against the expectations the first and the second canonical traits were found very highly significant ($P < 0.001$) and the fourth was slightly significant ($P < 0.05$) for the contrasts between the AB and BB pattern. This result showed that Pit-1 could have more than one action. The first canonical trait is more specifically linked to angularity. The last trait reflected the equilibrium between milk and protein yields. In order to make these contrasts more understandable, table 7 gives the values of the contrasts and the standard errors expressed on the original scales. We observed that the backtransformed contrasts were very important for milk, fat and protein for the first canonical contrast. All were also positive with AB animals superior to BB animals. For the second canonical trait the AB were inferior for milk, fat and protein and superior for angularity. This indicates again that the influence of Pit-1 on angularity seems to be important, first through the link between yields and angularity, but also directly on angularity with a slightly negative influence on yields. Canonical trait three did not show significant contrasts and canonical trait four, despite being significant,

explained only very little of the total variance. After grouping all the significant canonical traits together, we observed higher grouped contrasts as in the single-trait situation. This was especially clear for fat yield and angularity, but also for milk and protein. The reason seems to be that the multiple-trait contrasts include information from the correlated traits, especially for fat and angularity this could explain the differences. Standard errors of contrasts did not increase in an important way, they were even reduced for milk and fat yields.

10 **TABLE 6.** Linear contrasts (C) and standard errors (SE) between the three Pit-1 patterns for the four canonical traits observed on 89 Holstein-Friesian bulls.

Canonical	Contrast					
	AA-AB ¹		AA-BB ¹		AB-BB	
trait	C	SE	C	SE	C	SE
1	-0.093	0.098	0.023	0.093	0.116***	0.017
2	0.003	0.102	-0.032	0.052	-0.035***	0.009
3	-0.021	0.038	-0.016	0.037	0.005	0.007
4 ²	0.005	0.021	-0.004	0.021	-0.009*	0.004

¹ Only 2.2% of the animal were AA, therefore all results comparing this pattern are preliminary.

20

² Eigenvalue associated with canonical trait 4 was very low, therefore the results should be interpreted as non-significant.

* P < 0.05

*** P < 0.001

25

25

TABLE 7. Linear contrast (C) and standard error of contrast (SE) between AB and BB obtained by backtransformation on 89 Holstein-Friesian bulls.

5	Trait	Canonical trait									
		1***		2***		3		4*		All significant ¹	
		C	SE	C	SE	C	SE	C	SE	C	SE
	Milk yield	289	44	-57	15	9	13	-27	11	205	48
	Fat yield	12.6	1.9	-0.3	0.1	-0.9	1.2	-0.1	0.1	12.1	1.9
	Protein yield	8.6	1.3	-1.2	0.3	0.2	0.3	0.9	0.4	8.3	1.4
	Angularity	1.126	0.169	0.782	0.211	0.223	0.032	-0.010	0.004	1.897	0.27
											1

¹ Combined linear contrast using the three significant canonical traits.

* P < 0.05

10 *** P < 0.001

EXAMPLE B

Sequencing of the Pit-1 gene and characterization of a mutation.

This method generates separate populations of radiolabeled oligonucleotides that begin from a fixed point and terminate randomly at a fixed residue or combination of residues. Because every base in the DNA has an equal chance of being a variable terminus, each population consists of a mixture of oligonucleotides whose lengths are determined by the location of a particular base along the length of the original DNA. These populations of oligonucleotides are then resolved by electrophoresis under conditions that can discriminate between individual DNAs that differ in length by as little as one nucleotide. When the populations are loaded into adjacent lanes of a sequencing gel, the order of nucleotides along the DNA can be read directly from an autoradiographic image of the gel.

25

26

Reference: Sanger, F., S. Nicklen, and A.R. Coulson. 1977, DNA sequencing with chain-terminating inhibitors, *Proc. Natl. Acad. Sci.* 74:5463.

EXAMPLE C

5 Detection experiments using the primers.

1) Ligase Chain Reaction.

Ligase chain reaction (LCR), employing just oligonucleotide probes and DNA ligase, is capable of detecting approximately 1000 copies of a specific target DNA sequence in the presence of a vast excess of other DNA
 10 sequence information. Since the first description in 1989 (Backman and Wang, 1989, European Patent Application No. 0 320 308; Royer et al., 1989, European Patent Application No. 0 324 616; Wallace, 1989, European Patent Application No. 0 336 731; Wu and Wallace, 1989, *Genomics* 4:560-569; Orgel, 1989; Richards and Jones, 1989) LCR has been improved by the
 15 employment of a thermostable DNA ligase in conjunction with non-radioactive detection (Bond et al., 1990).

TABLE 8

20 Regression on the number of copies of the Pit-1 A allele (gene substitution effect) and on the presence of AB (dominance effect) observed for 455 valid records (lactation length 250-730 days) of 174 Cana cows.

Trait	Effect	Regression coefficient	SE	Pr > T
Milk yield	A allele	128	333	0.70
	Dominance (AB)	-136	378	0.72
Fat yield	A allele	1.40	14.27	0.92
	Dominance (AB)	-1.03	16.19	0.95
Protein yield	A allele	4.66	10.33	0.65
	Dominance (AB)	-5.74	11.72	0.62

2) FLP at the Pit-1 gene using HinfI restriction enzyme was revealed by
 25 PCR analysis adapted from Woolard et al., supra.

27

Using the method described by Sanger et al., supra, we have identified the point mutation at the nucleotide 1178 (a versus g) associated to the reproted RFLP. Also, a new PCR method without HinfI restriction enzyme and using primers that overlap the mutation has been developed.

5

Polymerase Chain Reaction Method

The RFLP at the Pit-1 gene was revealed by the polymerase chain reaction (PCR). Briefly, two PCR primers that overlap the mutation (primer AA = 5'-CAGAGAGAAAAACGGGTGAAGACAAGCATA-3' and primer BB = 5'-
10 CAGAGAGAAAAACGGGTGAAGACAAGCATG-3') were used in association with a third primer (primer B = 5'-
GACAGGGAAAGTGATATAGAAAGGGAGATAGA-3') to amplify a 360-bp fragment from the genomic DNA in a 50- μ l reaction volumes containing 2 mM MgCl₂. Conditions were 95°C for 3 min, followed by 35 cycles of 95°C for 1
15 min, 65.2°C for 1 min, and 72°C for 1 min. The final step was 72°C for 10 min. The PCR products were electrophoresed on 2% agarose gels with 1 μ g/ml of ethidium bromide (Figure 3).

28

CONCLUSIONS

Two alleles were distinguished for the Pit-1 gene, the growth hormone factor-1/pituitary-specific transcription factor responsible for the activation of prolactin and GH gene expression, using a restriction site recognized by *Hinfl*. Two allele were observed, A not digested and B showing this site. The AA pattern was less frequent than the AB or BB pattern. The significant superiority of the Pit-1 AB pattern or the AA pattern over BB was observed for milk, protein and angularity. This indicates that the heterozygote animals have higher productions and greater dairyness. The fat percentage was found to be lower for AB than for BB animals, a result that results from higher milk by near constant fat yield.

These results show a single action of Pit-1. But, by using a canonical transformation approach it was observed that at least two different actions of Pit-1; one on yields and angularity and another only on angularity. These results can be explain that Pit-1 has more than one role through the activation of prolactin and the GH gene expression. A first role is influencing milk, protein (and fat) yields, a second role is linked to the muscular development of the animals, meaning the presence of AB reducing the muscularity through an improvement of angularity.

Interesting enough, these findings show the usefulness of the canonical transformation to distinguish between effects on related traits. The association of Pit-1 polymorphism and milk traits in dairy cattle was shown on the original, but also on a transformed scale. Relationships were less important for conformation traits, except angularity, a trait that is related to milk yield. Again canonical transformation showed that effects on angularity were only partially a direct consequence of influence of Pit-1 on milk traits.

Identification of a specific mutation in the Pit-1 gene further allows the rapid and sensitive method to be carried out to distinguish between the various alleles and corresponding traits.

29

What we Claim is:

1. A genetic marker used to distinguish amongst animals a trait for milk producing capabilities or meat producing capabilities said genetic marker comprising a mutation in a fragment of a Pit-1 gene wherein three patterns of alleles are observed in which one of said allele patterns was fully mutated and being indicative of a trait of muscularity in said animal, while the two other allelic patterns, one being mutated and non/mutated, the other being non-mutated/non-mutated being indicative of a milk producing trait in said animal.
2. The genetic marker according to claim 1, wherein the mutation is a transition from A to G in the 1178 position of the sequence of Figure 2.
3. The genetic marker according to Claim 1 or Claim 2 wherein, the observation is via an electrophoretic gel.
4. The genetic marker according to any one of claims 1 to 3 ,wherein the muscularity trait has the BB allelic frequency.
5. The genetic marker according to any one of Claims 1 to 3, wherein the milk-producing trait has the AB allelic frequency or the AA allelic frequency.
6. A method for detecting certain traits in an animal, said method comprising the steps of:
 - (1) isolating genomic DNA from an animal;
 - (2) optionally isolating a fragment from said genomic DNA comprising a fragment of the Pit-1 gene;
 - (3) detecting a mutation in the Pit-1 gene; and
 - (4) analyzing said mutation to determine traits in said animal wherein upon analysis traits of muscularity and fat are distinguished from milk producing traits in said animal.
7. The method according to claim 6, wherein in step (3), detecting the mutation in the Pit-1 gene is performed using a restriction endonuclease.
8. The method according to Claim 7, wherein said restriction enzymes is HinfI.

9. The method according to Claim 6 or Claim 7, wherein said fragment of the Pit-1 gene is isolated using PCR primers.

10. The method according to Claim 9, wherein said PCR primers are:

5 5'-AAACCATCATCTCCCTTCTT-3' (SEQ ID NO:1)

5'AATGTACAATGTGCCTTCTGAG-3' (SEQ ID NO 2).

11. The method according to any one of Claim 7, wherein the detection step further comprises after digestion, analysis via RFLP or CFLP or SSCP or DGGE.

10 12. The method according to claim 6, wherein in step (3), detecting the mutation in the Pit-1 gene is performed using primer(s) which overlap(s) the mutation in said Pit-1 gene.

13. The method according to claim 12, wherein PCR primers are the following:

15 ca gagagaaaaa cgggtgaaga caagcat a (Pit-1 AA)

gac agggaaagtg atatagaaag ggagataga (Pit-1 B)

for the AA genotype, characteristic of milk producing capabilities, and

ca gagagaaaaa cgggtgaaga caagcat g (Pit-1 BB)

gac agggaaagtg atatagaaag ggagataga (Pit-1 B)

20 for the BB genotype, characteristic of meat producing capabilities.

14. A mutated Pit-1 gene comprising a mutation on position 1178.

15. A probe for detecting the milk or meat capabilities which comprises the nucleotide on position 1178 of the sequence of Figure 2.




FIGURE 1

2/4

"MSCQPFTSTDTFIPLNSESSATLPLIMHPSAAECLPVSNHATNV
MSTATGLHYSVPFCHYGNQSSTYGVMAGSLTPCLYKFPDHTLSHGFPMMHQPLLED
TAADFKQELRRKSKLVEEPIDMDSPEIRELEKFANEFKVRRIKLGYTQTNVGEALAAV
HGSEFSQTTICRFENLQLSFKNACKLKAILSKWLEEAQVVGALYNEKVGANERKRKR
TTISIAAKDALERHFGEQNKPSSEILRMAEELNLEKEVVRVWFCNRRQREKRVKTSL

1 gcaataactg igatttgaag ciaaccaaataaactaatcttattttggc tggagaagag

61 aaaggaatga aagtagaaac actcgctatt acacatagga gacccatct gaattcgaga

121 tgcctctag aaatagtaaa taaactctga ttcaggcttg tcttcacccg tttttctc

181 tgcctcggat acaaaaccaa accctacca ctctttctc caggtttagt tcttcagcca

241 tccgcaggat ctctgagag gaaggcttat tctgtctcc aaagtgctc tccagggcgt

301 cttagcagc aatactgatt gttgtctcc gtttctatc tttgtggga atgagtgcc

361 aacctttac ttcgactgat acctttatac ctctgaattc tgagtctct gcaactctg

421 ctctgataat gcatccagc gctgcggagt gcctaccggt ctccaaccac gccaccaacg

481 tgatgccac agcaacagga ctctattatt ctgttcttt ctgtcattat ggaaccagt

541 catcgaccta tggcgtgatg gcaggagct taacccttg tctttataag tttctgacc

601 acacgttgag tcatggtttt cctccatgc atcagcctct ccttcagag gacccactg

661 ccgctgatit caagcaggag ctaggcgga aaagcaaatt ggttgaagag ccaatagaca

721 tggattctcc agaaatccga gaacttgaaa agtttgccaa tgagtttaaa gtgagaagaa

FIGURE 2/1

3/4

781 ttaagctagg atacaccag acaaatgtg ggaagctci ggcagcttg catggctctg

841 aattcagica aacaactatc tggcatttg aaaacctgca gctcagcttc aaaaatgcat

901 gcaactaaa agcaatatta tccaaatggc tggaggaagc cgagcaagta ggagctttat

961 acaatgagaa agttgggca aatgaagaa aaaggaaacg gagaacaaca atcagtattg

1021 ctgctaaaaga cgcgctggag agacacttg gagaacagaa taagccttc tctcaggaga

1081 tcctgcggat ggctgaagaa ctaacctgg agaaagaagt ggtgagggtt tggttttgta

Hinf-I

1141 accgaaggca gagagaaaaa cgggtgaaga caagcct |aaa tc |agagtta ttactattt
 ca gagagaaaaa cgggtgaaga caagcat a (pit-1 AA)
 ca gagagaaaaa cgggtgaaga caagcat g (pit-1 BB)

1201 ctaaggagca tctgaatgc agataggctc tctattgtg taatagcgt tctactttc

1261 atctctct cttctcagcc aaaatagaaa ttggtattt ggtagcnnn aaaaatcaca

1321 tcagtaattt ttgncagaag ttttcttt ctactttaaa aataaataca atttaaat

1381 tgtgatgaa ntattctcag aaggannnnn tcaatgtaca nttaagcca aagactaata

1441 ggattaaaac aatgattctg t ccct t tcac tatatct t tc cctctatct tccnggaat
 gac agggaaagt atataagaaag ggagataga (pit-1-B)

1501 tc

FIGURE 2/2

1 2 | 3 4 | 5 6




FIGURE 3



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A3	(11) International Publication Number: WO 98/03677 (43) International Publication Date: 29 January 1998 (29.01.98)
(21) International Application Number: PCT/EP97/03939 (22) International Filing Date: 22 July 1997 (22.07.97) (30) Priority Data: 96401634.9 22 July 1996 (22.07.96) EP (34) Countries for which the regional or international application was filed: GB et al. (71) Applicant (for all designated States except US): BYOCORE LIMITED [GB/-]; P.O. Box 3175, Road Town, Tortola (VG). (72) Inventors; and (75) Inventors/Applicants (for US only): RENAVILLE, Robert [BE/BE]; Rue de Moha 18, B-5030 Gembloux (BE). PORTETELLE, Daniel [BE/BE]; Rue de la Motte 11, B-5081 Meux la Bruyère (BE). (74) Agents: GUTMANN, Ernest et al.; Ernest Gutmann-Yves Plasseraud S.A., 3, rue Chauveau-Lagarde, F-75008 Paris (FR).	(81) Designated States: CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 23 April 1998 (23.04.98)	
(54) Title: PIT-1 GENE POLYMORPHISM AND TRAIT SELECTION IN ANIMALS		
(57) Abstract The invention concerns a genetic marker used to distinguish amongst animals a trait for milk producing capabilities or meat producing capabilities said genetic marker comprising a mutation in a fragment of a Pit-1 gene wherein three patterns of alleles are observed in which one of said allele patterns was fully mutated and being indicative of a trait of muscularity in said animal, while the two other allelic patterns, one being mutated and non/mutated, the other being non-mutated/non-mutated being indicative of a milk producing trait in said animal.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 97/03939

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	YU ET AL.: "Association of PIT1 polymorphisms with growth" JOURNAL OF ANIMAL SCIENCE, vol. 73, no. 5, May 1995, CHAMPAIGN, ILL., pages 1282-1288, XP000197178 cited in the application see the whole document ---	1,3-12
Y	WOOLLARD ET AL.: "Rapid communication: HinfI polymorphism at the bovine PIT1 locus" JOURNAL OF ANIMAL SCIENCE, vol. 72, no. 12, December 1994, CHAMPAIGN, ILL., page 3267 XP000197179 cited in the application see the whole document ---	1,3-12
	-/--	

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

12 February 1998

Date of mailing of the international search report

03.03.98.

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ceder, O

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/EP 97/03939

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BODNER ET AL.: "The pituitary transcription factor GHF-1 is a homeobox containing protein" CELL, vol. 55, 1988, pages 505-518, XP002054526 ---	2,14
A	YU ET AL.: "Expression pattern, genomic cloning and RFLP analyses of the swine PIT-1 gene" ANIMAL GENETICS, vol. 25, no. 4, 1994, pages 229-233, XP002054527 ---	2,14
A	JP 06 090 762 A (EIKEN CHEM CO) 5 April 1994 see sequences ---	2,14
L	PATENT ABSTRACTS OF JAPAN vol. 18, no. 351 (C-1220), 4 July 1994 & JP 06 090762 A (EIKEN CHEM CO), 5 April 1994, english abstract of JP6090762 see abstract ---	2,14
A	WO 90 10714 A (WISCONSIN ALUMINI RESEARCH FOUNDATION) 20 September 1990 see abstract ---	1,3,6,11
A	PROSSER J: "DETECTING SINGLE-BASE MUTATIONS" TRENDS IN BIOTECHNOLOGY, vol. 11, June 1993, pages 238-246, XP000601507 cited in the application ---	
A	MEJDELL CM ET AL: "Associations of bovine lymphocyte antigens with milk and meat production traits of Norwegian cattle." J DAIRY SCI, DEC 1993, 76 (12) P3778-84, UNITED STATES, XP000647803 -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 97/03939

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 6090762 A	05-04-94	NONE	

WO 9010714 A	20-09-90	US 5041371 A	20-08-91
		EP 0424480 A	02-05-91
		JP 3504805 T	24-10-91
